



## KCNQ4 gene

potassium voltage-gated channel subfamily Q member 4

### Normal Function

The *KCNQ4* gene provides instructions for making a protein that is part of a family of potassium channels. These channels transport positively charged potassium atoms (potassium ions) between neighboring cells. The channels play a key role in the ability of cells to generate and transmit electrical signals. The specific function of a potassium channel depends on its protein components and its location in the body. Potassium channels made with the *KCNQ4* protein are found in certain cells of the inner ear and along part of the nerve pathway from the ear to the brain (the auditory pathway). To a lesser extent, *KCNQ4* potassium channels are also found in the heart and some other muscles.

Because *KCNQ4* potassium channels are present in the inner ear and auditory pathway, researchers have focused on their role in hearing. Hearing requires the conversion of sound waves to electrical nerve signals, which are then transmitted to the brain. This conversion involves many processes, including maintenance of the proper levels of potassium ions in the inner ear. *KCNQ4* channels help to maintain these levels, playing a critical role in the efficient transmission of electrical nerve signals from the inner ear to the brain.

### Health Conditions Related to Genetic Changes

#### nonsyndromic hearing loss


Several *KCNQ4* gene mutations have been reported in individuals with nonsyndromic hearing loss, which is loss of hearing that is not associated with other signs and symptoms. Mutations in this gene cause a form of nonsyndromic hearing loss called DFNA2. This form of hearing loss generally begins after a child learns to speak (postlingual) and particularly affects the ability to hear high-frequency sounds. DFNA2 is described as progressive, which means it becomes more severe over time.

Most *KCNQ4* gene mutations change one of the building blocks (amino acids) used to make the *KCNQ4* protein. Some mutations prevent the channel from reaching the cell membrane, where it is needed to transport potassium ions. Other mutations lead to the formation of abnormal channels that cannot transport these ions effectively. The loss of functional *KCNQ4* channels appears to cause a buildup of potassium ions in certain cells of the inner ear, which damages those cells and leads to progressive hearing loss in people with DFNA2.

## Chromosomal Location

Cytogenetic Location: 1p34.2, which is the short (p) arm of chromosome 1 at position 34.2

Molecular Location: base pairs 40,784,012 to 40,840,457 on chromosome 1 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

## Other Names for This Gene

- DFNA2
- KCNQ4\_HUMAN
- KQT-like 4
- KV7.4
- potassium channel, voltage gated KQT-like subfamily Q, member 4
- potassium voltage-gated channel, KQT-like subfamily, member 4

## Additional Information & Resources

### Educational Resources

- Biochemistry (fifth edition, 2002): Specific Channels Can Rapidly Transport Ions Across Membranes  
<https://www.ncbi.nlm.nih.gov/books/NBK22509/>
- Neuromuscular Disease Center, Washington University: KCNQ family of potassium channels  
<http://neuromuscular.wustl.edu/mother/chan.html#kcnqfam>

### GeneReviews

- Deafness and Hereditary Hearing Loss Overview  
<https://www.ncbi.nlm.nih.gov/books/NBK1434>
- DFNA2 Nonsyndromic Hearing Loss  
<https://www.ncbi.nlm.nih.gov/books/NBK1209>

### Scientific Articles on PubMed

- PubMed  
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28KCNQ4%5BTIAB%5D%29+OR+%28DFNA2%5BTIAB%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

### OMIM

- POTASSIUM CHANNEL, VOLTAGE-GATED, KQT-LIKE SUBFAMILY, MEMBER 4  
<http://omim.org/entry/603537>

### Research Resources

- ClinVar  
<https://www.ncbi.nlm.nih.gov/clinvar?term=KCNQ4%5Bgene%5D>
- Hereditary Hearing Loss Homepage  
<http://hereditaryhearingloss.org>
- HGNC Gene Family: Deafness associated genes  
<http://www.genenames.org/cgi-bin/genefamilies/set/1152>
- HGNC Gene Family: Potassium voltage-gated channels  
<http://www.genenames.org/cgi-bin/genefamilies/set/274>
- HGNC Gene Symbol Report  
[http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?q=data/hgnc\\_data.php&hgnc\\_id=6298](http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=6298)
- NCBI Gene  
<https://www.ncbi.nlm.nih.gov/gene/9132>
- UniProt  
<http://www.uniprot.org/uniprot/P56696>

### **Sources for This Summary**

- Coucke PJ, Van Hauwe P, Kelley PM, Kunst H, Schatteman I, Van Velzen D, Meyers J, Ensink RJ, Verstreken M, Declau F, Marres H, Kastury K, Bhasin S, McGuirt WT, Smith RJ, Cremers CW, Van de Heyning P, Willems PJ, Smith SD, Van Camp G. Mutations in the KCNQ4 gene are responsible for autosomal dominant deafness in four DFNA2 families. *Hum Mol Genet.* 1999 Jul;8(7):1321-8.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/10369879>
- Gao Y, Yechikov S, Vázquez AE, Chen D, Nie L. Impaired surface expression and conductance of the KCNQ4 channel lead to sensorineural hearing loss. *J Cell Mol Med.* 2013 Jul;17(7):889-900.  
doi: 10.1111/jcmm.12080. Epub 2013 Jun 11.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/23750663>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3729637/>
- GeneReview: DFNA2 Nonsyndromic Hearing Loss  
<https://www.ncbi.nlm.nih.gov/books/NBK1209>

- Kim HJ, Lv P, Sihn CR, Yamoah EN. Cellular and molecular mechanisms of autosomal dominant form of progressive hearing loss, DFNA2. J Biol Chem. 2011 Jan 14;286(2):1517-27. doi: 10.1074/jbc.M110.179010. Epub 2010 Oct 21.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/20966080>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3020760/>
- Kubisch C, Schroeder BC, Friedrich T, Lütjohann B, El-Amraoui A, Marlin S, Petit C, Jentsch TJ. KCNQ4, a novel potassium channel expressed in sensory outer hair cells, is mutated in dominant deafness. Cell. 1999 Feb 5;96(3):437-46.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/10025409>
- Naito T, Nishio SY, Iwasa Y, Yano T, Kumakawa K, Abe S, Ishikawa K, Kojima H, Namba A, Oshikawa C, Usami S. Comprehensive genetic screening of KCNQ4 in a large autosomal dominant nonsyndromic hearing loss cohort: genotype-phenotype correlations and a founder mutation. PLoS One. 2013 May 23;8(5):e63231. doi: 10.1371/journal.pone.0063231. Print 2013.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/23717403>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3662675/>
- OMIM: POTASSIUM CHANNEL, VOLTAGE-GATED, KQT-LIKE SUBFAMILY, MEMBER 4  
<http://omim.org/entry/603537>

---

Reprinted from Genetics Home Reference:  
<https://ghr.nlm.nih.gov/gene/KCNQ4>

Reviewed: February 2016  
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications  
U.S. National Library of Medicine  
National Institutes of Health  
Department of Health & Human Services